


High-redshift Quasar Studies in the TMT era

Masafusa Onoue (Kavli IPMU / KIAA)

SUBARU USERS MEETING FY2024, 30 JAN 2025

IPMU INSTITUTE FOR THE PHYSICS AND
MATHEMATICS OF THE UNIVERSE







Supermassive Black Hole

- Residing at the galaxy center
- Mass: $\log M_{\text{BH}} / M_{\text{sun}} = 5 - 10$
- Observable when mass accretes onto BH (AGN/quasar)
- $10^9 M_{\text{sun}}$ BH observed within 1 Gyr of the Universe

AGN discovery (2000s-)




ground-based wide-field surveys
(SDSS, Pan-Starrs1)




UV-luminous unobscured quasars
($\log L_{\text{bol}} [\text{erg s}^{-1}] = 47, \log M_{\text{BH}} / M_{\text{sun}} > 9$)

AGN discovery (2020s)




space-based surveys
(JWST)







Optical-selected unobscured+obscured AGN
 $(\log L_{\text{bol}} [\text{erg s}^{-1}] = 44 - 45, \log M_{\text{BH}} / M_{\text{sun}} = 6-7)$


$z > 10$ Galaxies Hosting AGN



- $z=10.1$ X-ray AGN (Goulding+23; Bogdan+24)
- GN-z11 ($z=10.6$; Maiolino+24)




Subaru's role?




→ **Intermediate-luminosity quasars**
(more representative BHs than SDSS, more matured than JWST)

HSC-SSP's constraints on $z > 4$ quasar abundance





SDSS limit




- ◆ $z=4-7$ Luminosity function constrained down to faint end
- ◆ HSC is dominating the quasar discovery at intermediate-luminosity range (until LSST)

JWST follow-up of HSC z=6 quasars





Host stellar emission can be well recovered with JWST ($\log M^*/M_{sun} \sim 10.5$)
& secure M_{BH} estimate with Hβ

Full Cy1 Sample (F356W)



Detection rate: 10 / 12, $\log M_*/M_{\text{sun}} \sim 10.5$ (host fraction $\sim 10\text{-}40\%$)

Quiescent Quasar Host



◆ **NIRCam imaging + NIRSpect G395M shows**

- clear Balmer break (F250M+F300M)
- no spatially extended H α (F480M)
- age = 240^{+30}_{-40} Myr ($z_{\text{form}}=8.2\pm0.4$)
- $\text{sSFR}_{100\text{Myr}} < 0.1 \text{ Gyr}^{-1}$
- $\sigma_* < 180 \pm 50 \text{ km s}^{-1}$ (from Balmer lines)


Limitation in JWST studies

- ◆ **Spectroscopic confirmation is limited**
($M_{UV} > 28$ mag or $\log M_{BH}/M_{sun} < 6$ candidates are too faint for NIRSpec)
- ◆ **Does PSF really represent quasar emission?** There could be possible contribution of *unresolved stellar emission* to PSF (, in which case M_* is underestimated)
- ◆ $M_{BH} - \sigma_*$ relation can be directly compared with the local relation, but **σ_* measurements for quasar hosts are challenging** (success rate for HSC quasars: 2/12)
- ◆ **Sample size is limited** ($O \sim 10$ at $z \sim 6$; $O \sim 100$ at $z \sim 2$; Tanaka+24)
 - ➡ Need a larger sample to constrain *scatter* of M_* / M_{BH} , but JWST observing time is too expensive
 - ➡ M_{BH} estimate in NIR is also expensive with 8m-class telescopes (> 5 hrs / target at $M_{UV} > 23$)



TOWARD TMT ERA

TMT's sensitivity



At 1-2 μm , AO-assisted sensitivity is comparable to JWST at 1-2 μm
Better in spectroscopy (at wavelengths not affected by OH sky)



TMT Science cases



◆ Systematic TMT+AO follow-up of HSC/PFS faint quasars (at $z \leq 4$)

- Host stellar light detection for M_* (to cover $\lambda_{\text{rest}} > 4000\text{\AA}$)
- M_{BH} estimate through rest UV lines, e.g., MgII 2798, CIV 1400 (especially at >23 mag)
- Deep NIR spectroscopy to detect host stellar absorption ($M_{\text{BH}} - \sigma_*$)

* **A large sample ($N > 100$)** enables us to constrain the BH demographics in the M_* / σ_* vs M_{BH} plane at high redshift (*BH first and/or galaxy first*)



TMT Science cases



Other science cases


✦ high-SNR spectral follow-up to characterize weak emissions

- Broad-line-region metallicity (e.g., H β 1640 vs C IV 1400)
- Fe II / Mg II ($\sim$ [Fe / α] on the nuclear scale)

✦ IFU follow up to trace AGN feedback: to be compared with ALMA

✦ Confirmation of JWST/Euclid/Roman-selected faintest AGN candidates

- M_{BH} distribution; seed BH signature (e.g., O I λ 1304, C II λ 2326; Inayoshi, MO+22)



Summary

- ◆ **Subaru has played a significant role in high- z quasar observations, bridging the gap between SDSS and JWST discovery space**
 - ◆ **JWST follow-up observations of HSC quasars have revealed host stellar emission at $z=6$, thanks to their relatively faint quasar luminosity.** Characterization of M^* vs M_{BH} distribution is ongoing.
 - ◆ **TMT follow-up observations of Subaru(HSC/PFS)-selected quasars will reveal the demographics of high- z SMBHs,** overcoming some limitations in the JWST studies and possibly revealing young seed BHs
-