


The Cosmic Himalayas:
**The Highest QSO Density Peak at Cosmic Noon
Bridging AGNs, Galaxies, and IGM**

Yongming Liang

NAOJ → ICRR Fellow, UTokyo

In collaborations with:


M. Ouchi (NAOJ/ICRR), N. Kashikawa (Tokyo), Z. Cai (Tsinghua), K. Nagamine (Osaka), S. Cantalupo (UniMIB), H. Yajima (Tsukuba), X. Fan (Arizona), M. Tanaka (NAOJ), R. Shimakawa, M. Onoue (Waseda), et al.




Background: Galaxy Evolution and AGN

◆ Galaxy Evolution with QSO/AGN in Theory

(Hopkins+2008)




◆ AGNs Triggered by Mergers?



Evidence is found at $z < \sim 1$ (e.g., Goulding+2019)


BUT: How about higher- z ? Type-1 QSO?

◆ QSOs Trace Protoclusters?



Background: AGNs in Overdensities


- ◆ **AGN Fraction in (proto-)clusters:** increasing towards high- z and peak at $z \sim 2$?



(Vito+2024)


3. An Extreme QSO Overdensity

◆ The Most Clustered SDSS/eBOSS QSOs@ $z=2.2$ Over 10,000 deg²



An Extreme QSO Overdensity

◆ The Most Clustered SDSS QSOs@z=2.2 Over 10,000 deg²





IGM Tomography Methodology

◆ 3D HI Tomography from SDSS/eBOSS QSOs

- Calculate δ_F for each lines-of-sight
 - Intrinsic continuum by MF-PCA.
 - Transmission at each pixel
 - Ly α transmission contrast δ_F compared to cosmic mean.



$$\delta_F = F_{\text{Ly}\alpha} / \langle F(z) \rangle_{\text{cos}} - 1$$



IGM Tomography Result

◆ 3D HI Tomography from SDSS/eBOSS QSOs: Collapsed Map


- Distinct ionizing stages between the two LAE nodes.
- QSOs centered at the transitioning regions.



Cosmic Himalayas: IGM Tomography Result

◆ Possible Ionizing Sources in the Ionizing Boundary

- Contributions from luminous quasars
- A significant correlation between the local HI transparency and $L_{\text{bol,QSO}}$



Cosmic Himalayas: IGM Tomography Result


◆ 3D HI Tomography from SDSS/eBOSS QSOs: Slicing Map

- Photoionization contributed in the QSO proximity zone:

$$\Gamma_{\text{HI}, i}^{\text{QSO}} = 8.8 \times 10^{-7} \xi \left(\frac{r_i}{10 \text{ kpc}} \right)^{-2}$$

- Photoionization radius that QSO > UVB

$$\Gamma_{\text{HI}, i}^{\text{QSO}} > \Gamma_{\text{HI}}^{\text{UVB}} \sim 10^{-12} \text{ s}^{-1}$$





Liang et al. 2025, ApJ, 986, 60


Cosmic Himalayas: How are the QSOs triggered?

◆ Possible Interpretations of the Quasar Overdensity

- An extreme protocluster w/ rich dust or heated gas, suppressing LAEs?
- A progenitor of cluster merger or intersection of filaments at $z=2.2$?



Clowe+2004




Chiang+2013

Cosmic Himalayas: How are the QSOs triggered?

◆ Possible Interpretations of the Quasar Overdensity


- The Dusty Population Revealed by JCMT/SCUBA-2 850 μ m down to **3 mJy**



Cosmic Himalayas: How are the QSOs triggered?

◆ Galaxy evolution along a filament where dust play a crucial role

- JCMT/SCUBA-2 850um maps also indicate a transitioning nature?



Future Perspective with Subaru/PFS


◆ PFS observation: evolution along a filament of diverse galaxies and fined IGM structures

- **Spec-z & galactic properties of photometric galaxies:**
 - LAEs (#450) + LBG/BX Gal. + HAEs/BzK + SMGs + X-ray/Radio sources...
(+u image) (+NB, K image) (+JCMT) (+Chandra/MeerKAT)
- **LOS sampling from background QSOs to LBGs:** $N_{\text{LoS}} = 23 \rightarrow \text{O}(1000)$
 - $d_{\perp} \sim 20 \text{ cMpc} \rightarrow d_{\perp} \sim 4\text{-}2 \text{ cMpc}$ @ 6-12 hrs (ref. SSP GE survey; Greene+2022)


u+K Imaging finished in
2024B +2025B:

1. CFHT MegaPrime (u)
2. Subaru MOIRCS (K, NB)
3. Blanco-4m NEWFIRM (K, NB)

MeerKAT L-band finished
in 2024B



With PFS




Future Perspective with X-ray Observation


◆ Chandra X-ray observation: SMBH growth and QSO triggering associated with the ongoing accretion disk

340 ks to $L_{2-10\text{keV}} = 10^{43.7} \text{ erg/s}$


Host Properties
 $N_H, \lambda_{\text{acc}}, \text{XLF}$



AGN Fraction vs. Stellar Mass Limit



AGN Accretion vs. Distance to QSOs



(Travascio+2024)

○ Chandra Cycle 27 Approved Targets

https://cxc.cfa.harvard.edu/target_lists/cycle27/cycle27_approved_peer_targets_by_radec.html

01:44:04.1	-15:56:14.9	tau Ceti	HRC-S	NONE	60.00	27200138	Mélis	University of California, San Diego (UCSD)
01:46:35.7	49:14:43.1	ZTF 0146	ACIS-S	NONE	50.00	27300224	Cunningham	Harvard University
01:48:37.6	06:00:20.0	SDSS J014837.63+0600	ACIS-I	NONE	750.00	27700099	Eilers	Massachusetts Institute of Technology (MIT)
01:57:50.0	62:12:26.7	PSR B0154+61	ACIS-S	NONE	60.00	27500298	Ng	McGill University
02:08:48.0	01:00:00.0	BOSS0210E	ACIS-I	NONE	170.00	27700042	Lepore	INAF/OA - Arcetri
02:10:00.0	00:50:24.0	CH Chandra01	ACIS-I	NONE	340.00	27700187	Liang	National Astronomy Observatory of Japan (NAOJ)
02:10:56.4	58:45:17.7	PSR J0210+5845	ACIS-S	NONE	60.00	27500298	Ng	McGill University
02:11:00.0	00:46:48.0	BOSS0210W	ACIS-I	NONE	170.00	27700042	Lepore	INAF/OA - Arcetri
02:42:41.3	-00:00:31.9	SN 2018ivc	ACIS-S	NONE	40.00	27500336	Chandra	NRAO/Charlottesville
03:14:26.1	-01:08:46.6	J0314-0108	ACIS-S	NONE	30.00	27620318	Osknova	Universitaet Potsdam
03:33:16.7	-26:53:46.3	SpARCS 0333	ACIS-S	NONE	280.00	27800080	Noble	Arizona State University (ASU)
03:41:41.4	52:57:28.3	1LHAASO J0343+5154u	ACIS-I	NONE	50.00	27910143	Mori	Columbia University
03:43:05.9	52:47:48.4	1LHAASO J0343+5154u	ACIS-I	NONE	50.00	27910143	Mori	Columbia University
03:44:02.9	-06:53:00.6	QSO J0344-0653	ACIS-S	NONE	60.00	27700175	Ighina	Smithsonian Astrophysical Observatory (SAO)

Summary

The Cosmic Himalayas:

An unprecedented quasar overdensity ($\delta_{\text{QSO}} \sim 17\sigma$ if Gaussian LSS) at $z = 2.2$, serving as a unique laboratory to study galaxy and black hole co-evolution at Cosmic Noon.

- ◆ QSOs located at ionization transition zones, jointly shaping IGM environments.
- ◆ QSO luminosity correlated with HI transparency, highlighting **quasar-driven ionization processes on tens of Mpc scales**.
- ◆ Spatially offset populations of LAEs, QSOs, and dusty SMGs, indicating diverse galaxy **evolutionary stages and feedback mechanisms**.

Subaru/PFS is of high necessity:

- ◆ The combination of **Wide FOV & Highly Multiplex Fibers & Large Telescope** is crucial to simultaneously take the spectroscopic data for diverse **foreground** galaxies and **background** LBGs in the ~ 100 -Mpc scale structures.



Liang et al. 2025,
ApJ, 986, 60

